



PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 11/21/2019
LAST REVIEW DATE: 11/18/2021
LAST CRITERIA REVISION DATE: 11/18/2021
ARCHIVE DATE:

NOURIANZ™ (istradefylline)

Coverage for services, procedures, medical devices and drugs are dependent upon benefit eligibility as outlined in the member's specific benefit plan. This Pharmacy Coverage Guideline must be read in its entirety to determine coverage eligibility, if any.

This Pharmacy Coverage Guideline provides information related to coverage determinations only and does not imply that a service or treatment is clinically appropriate or inappropriate. The provider and the member are responsible for all decisions regarding the appropriateness of care. Providers should provide BCBSAZ complete medical rationale when requesting any exceptions to these guidelines.

The section identified as "**Description**" defines or describes a service, procedure, medical device or drug and is in no way intended as a statement of medical necessity and/or coverage.

The section identified as "**Criteria**" defines criteria to determine whether a service, procedure, medical device or drug is considered medically necessary or experimental or investigational.

State or federal mandates, e.g., FEP program, may dictate that any drug, device or biological product approved by the U.S. Food and Drug Administration (FDA) may not be considered experimental or investigational and thus the drug, device or biological product may be assessed only on the basis of medical necessity.

Pharmacy Coverage Guidelines are subject to change as new information becomes available.

For purposes of this Pharmacy Coverage Guideline, the terms "experimental" and "investigational" are considered to be interchangeable.

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

This Pharmacy Coverage Guideline does not apply to FEP or other states' Blues Plans.

Information about medications that require precertification is available at www.azblue.com/pharmacy.

Some large (100+) benefit plan groups may customize certain benefits, including adding or deleting precertification requirements.

All applicable benefit plan provisions apply, e.g., waiting periods, limitations, exclusions, waivers and benefit maximums.

Precertification for medication(s) or product(s) indicated in this guideline requires completion of the [request form](#) in its entirety with the chart notes as documentation. **All requested data must be provided.** Once completed the form must be signed by the prescribing provider and faxed back to BCBSAZ Pharmacy Management at (602) 864-3126 or emailed to Pharmacyprecert@azblue.com. **Incomplete forms or forms without the chart notes will be returned.**



PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 11/21/2019
LAST REVIEW DATE: 11/18/2021
LAST CRITERIA REVISION DATE: 11/18/2021
ARCHIVE DATE:

NOURIANZ™ (istradefylline)

Criteria:

- **Criteria for initial therapy:** Nourianz (istradefylline) is considered *medically necessary* and will be approved when **ALL** of the following criteria are met:
1. Prescriber is a physician specializing in patient's diagnosis or is in consultation with a Neurologist
 2. Individual is 18 years of age or older
 3. A confirmed diagnosis of Parkinson's disease (PD) in an individual experiencing "off" episodes and **ALL** of the following:
 - a. Parkinson's disease is Hoehn and Yahr Stage 2 or greater
 - b. There is at least 2 hours of off time per day
 4. Use is as adjunctive treatment to levodopa/carbidopa
 5. Individual has failure, contraindication or intolerance to **ALL** of the following:
 - a. **One trial** of dopamine agonist: pramipexole, **or** ropinirole
 - b. **One trial** of monoamine oxidase inhibitor (MAO) B inhibitor: selegiline (capsule or tablet) **or** rasagiline mesylate tablet
 - c. **One trial** of catechol O-methylase inhibitor (COMT): entacapone **or** tolcapone
 6. Will not be used in severe renal impairment (CrCl < 15 mL/min) or end stage renal disease requiring dialysis
 7. Will not be used in severe hepatic impairment (Child-Pugh Class C)
 8. There are no significant interacting drugs

Initial approval duration: 6 months

- **Criteria for continuation of coverage (renewal request):** Nourianz (istradefylline) is considered *medically necessary* and will be approved when **ALL** of the following criteria are met:
1. Individual continues to be seen by a physician specializing in patient's diagnosis or is in consultation with a Neurologist
 2. Individual's condition responded while on therapy
 - a. Response is defined as **THREE** of the following:
 - i. No evidence of disease progression
 - ii. Functionality retained most activities of daily living
 - iii. Achieved and maintains a reduction in off time
 - iv. Achieved and maintains an increase in on time of at least 1 hour



PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 11/21/2019
LAST REVIEW DATE: 11/18/2021
LAST CRITERIA REVISION DATE: 11/18/2021
ARCHIVE DATE:

NOURIANZ™ (istradefylline)

3. Individual has been adherent with the medication
4. Individual has not developed any significant adverse drug effects that may exclude continued use
 - a. Significant adverse effect such as:
 - i. Emergent or worsening dyskinesia
 - ii. Emergent or worsening hallucinations or psychotic behaviors
 - iii. Impulse control or compulsive behaviors
5. Will not be used in severe renal impairment (CrCl < 15 mL/min) or end stage renal disease requiring dialysis
6. Will not be used in severe hepatic impairment (Child-Pugh Class C)
7. There are no significant interacting drugs

Renewal duration: 12 months

➤ Criteria for a request for non-FDA use or indication, treatment with dosing, frequency, or duration outside the FDA-approved dosing, frequency, and duration, refer to one of the following Pharmacy Coverage Guideline:

1. **Off-Label Use of a Non-cancer Medications**
2. **Off-Label Use of a Cancer Medication for the Treatment of Cancer without a Specific Coverage Guideline**

Description:

Nourianz (istradefylline) is an adenosine receptor antagonist indicated as adjunctive treatment to levodopa/carbidopa in adult patients with Parkinson's disease (PD) experiencing "off" episodes. The mechanism of action of istradefylline is unknown. In *in vitro* studies and in *in vivo* animal studies, istradefylline was demonstrated to be an adenosine A_{2A} receptor antagonist.

Patients with Parkinson disease (PD) who take levodopa chronically are increasingly likely to develop motor fluctuations and dyskinesia as the disease progresses and nigrostriatal dopaminergic neurons continue to degenerate and lose presynaptic dopamine storage capacity.

Dyskinesia consists of several types of abnormal involuntary movements, most often choreiform, brought on by levodopa or other dopaminergic agents. Dyskinesia is caused by overstimulation of dopamine receptors by the use of levodopa, but also by agents that either stimulate or enhance the effect of dopamine at the receptor (e.g., dopamine agonists, monoamine oxidase type B [MAO B] inhibitors, and catechol-O-methyl transferase [COMT] inhibitors).



**PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS**

**ORIGINAL EFFECTIVE DATE: 11/21/2019
LAST REVIEW DATE: 11/18/2021
LAST CRITERIA REVISION DATE: 11/18/2021
ARCHIVE DATE:**

NOURIANZ™ (istradefylline)

Motor fluctuations are alterations between periods of a positive response to medication (commonly referred to as "on" periods), and periods of reemergence of parkinsonian symptoms (commonly referred to as "off" periods). The transition from on to off can be sudden and unpredictable. Unpredictable off periods can also occur that have no relationship to timing of levodopa and the wearing off phenomena. Use of home diaries and direct observation can help determine any relationship of levodopa dosing to the off period. Many unpredictable off moments turn out to be end of levodopa dosing wearing off.

"Wearing off" is characterized by the recurrence of parkinsonian symptoms as the effect of levodopa diminishes near the end of the dose interval, usually three to four hours after a dose. Wearing off is often the first and most commonly encountered fluctuation. A majority of patients experience wearing off after five years of levodopa treatment; however, in a minority this phenomenon may occur earlier. Other motor fluctuations include freezing of gait that can lead to falls and ultimately loss of independence, acute akinesia, and failed or lack of an on response to levodopa.

Wearing off may be managed initially by increasing the dose of levodopa and/or interval adjustments and use of longer acting levodopa preparations. If levodopa adjustments for wearing off are not adequate or tolerated, the addition of an adjunctive therapy (e.g., dopamine agonist (pramipexole or ropinirole), COMT inhibitor, MAO B inhibitor, or istradefylline) to the levodopa regimen can help to reduce "off" time.

Definitions:

Hoehn and Yahr Scale and the Modified Hoehn and Yahr Scale:

Stage	Hoehn and Yahr Scale	Modified Hoehn and Yahr Scale
0	No signs of disease	No signs of disease
1	Unilateral involvement only, usually minimal or no functional disability	Unilateral involvement only
1.5	--	Unilateral and axial involvement
2	Bilateral or midline involvement without impairment of balance	Bilateral involvement without impairment of balance
2.5	--	Mild bilateral disease with recovery on pull test
3	Bilateral disease: mild to moderate disability with impaired postural reflexes; physically independent	Mild to moderate bilateral disease; some postural instability; physically independent
4	Severely disabling disease; still able to walk or stand unassisted	Severely disability; still able to walk or stand unassisted
5	Confined to bed or wheelchair unless aided	Wheelchair bound or bedridden unless aided
Hoehn M, Yahr M: Parkinsonism: onset, progression and mortality. Neurology 1967; 17 (5):427-442		

PHARMACY COVERAGE GUIDELINES
SECTION: DRUGS

ORIGINAL EFFECTIVE DATE: 11/21/2019
LAST REVIEW DATE: 11/18/2021
LAST CRITERIA REVISION DATE: 11/18/2021
ARCHIVE DATE:

NOURIANZ™ (istradefylline)

The Child-Pugh classification system:

The Child-Pugh classification is a scoring system used to determine the prognosis of individuals with cirrhosis. Scoring is based upon several factors: albumin, ascites, total bilirubin, prothrombin time, and encephalopathy, as follows:

	Score: 1 point	Score: 2 points	Score: 3 points
Serum Albumin (g/dL)	>3.5	3.0 - 3.5	<3.0
Serum Bilirubin (mg/dL)	<2.0	2.0 - 3.0	>3.0
Prothrombin time (seconds)	1 - 4	4 - 6	>6
Ascites	none	moderate	severe
Encephalopathy	none	mild	severe

The three classes and their scores are:

- **Class A** is score 5 – 6: Well compensated
- **Class B** is score 7 – 9: Significant functional compromise
- **Class C** is score >9: Decompensated disease

Resources:

Nourianz (istradefylline) product information, revised by Kyowa Kirin Inc 05-2020. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed September 06, 2021.

Liang TW, Tarsy D. Medical management of motor fluctuations and dyskinesia in Parkinson disease. In: UpToDate, Hurtig HI, Eichler AF (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Accessed July 26, 2021.