

PHARMACY COVERAGE GUIDELINE

RETEVMO™ (selpercatinib) oral

This Pharmacy Coverage Guideline (PCG):

- Provides information about the reasons, basis, and information sources we use for coverage decisions
- Is not an opinion that a drug (collectively “Service”) is clinically appropriate or inappropriate for a patient
- Is not a substitute for a provider’s judgment (Provider and patient are responsible for all decisions about appropriateness of care)
- Is subject to all provisions e.g. (benefit coverage, limits, and exclusions) in the member’s benefit plan; and
- Is subject to change as new information becomes available.

Scope

- This PCG applies to Commercial and Marketplace plans
- This PCG does not apply to the Federal Employee Program, Medicare Advantage, Medicaid or members of out-of-state Blue Cross and/or Blue Shield Plans

Instructions & Guidance

- To determine whether a member is eligible for the Service, read the entire PCG.
- This PCG is used for FDA approved indications including, but not limited to, a diagnosis and/or treatment with dosing, frequency, and duration.
- Use of a drug outside the FDA approved guidelines, refer to the appropriate Off-Label Use policy.
- The “Criteria” section outlines the factors and information we use to decide if the Service is medically necessary as defined in the Member’s benefit plan.
- The “Description” section describes the Service.
- The “Definition” section defines certain words, terms or items within the policy and may include tables and charts.
- The “Resources” section lists the information and materials we considered in developing this PCG
- **We do not accept patient use of samples as evidence of an initial course of treatment, justification for continuation of therapy, or evidence of adequate trial and failure.**
- Information about medications that require precertification is available at www.azblue.com/pharmacy. You must fully complete the [request form](#) and provide chart notes, lab workup and any other supporting documentation. The prescribing provider must sign the form. Fax the form to BCBSAZ Pharmacy Management at (602) 864-3126 or email it to Pharmacyprecert@azblue.com.

Criteria:

- **Criteria for initial therapy:** Retevmo (selpercatinib) is considered **medically necessary** and will be approved when **ALL** the following criteria are met:
 1. Prescriber is a physician specializing in the patient’s diagnosis or is in consultation with an Oncologist.
 2. Individual has a confirmed diagnosis of **ONE** of the following:
 - a. Adult patient (18 years of age or older) with metastatic *RET* fusion-positive non-small cell lung cancer (NSCLC)
 - b. Adult or pediatric patient 12 years of age or older with advanced or metastatic *RET*-mutant medullary thyroid cancer (MTC) who require systemic therapy

PHARMACY COVERAGE GUIDELINE

RETEVMO™ (selpercatinib) oral

- c. Adult or pediatric patient 12 years of age or older with advanced or metastatic *RET* fusion-positive thyroid cancer who require systemic therapy and who are radioactive iodine-refractory (if radioactive iodine is appropriate)
 - d. Other request for a specific oncologic direct treatment use that is found and listed in the National Comprehensive Cancer Network (NCCN) Guidelines with Categories of Evidence and Consensus of 1 and 2A
3. The individual has received and completed **ALL** the following **baseline tests** before initiation of treatment and with continued monitoring of the individual as clinically appropriate:
 - a. Adequately controlled blood pressure
 - b. EKG to assess QT interval
 - c. Potassium, magnesium, and calcium are within their normal ranges
 - d. TSH
 - e. Negative pregnancy test in a woman of childbearing potential
 - f. Eastern Cooperative Oncology Group (ECOG) Performance Status is ≤ 2
 4. Individual does not have active cardiovascular disease or a recent myocardial infarction.
 5. Individual does not have end-stage renal disease (eGFR less than 15 mL/min).
 6. Individual is not using strong and moderate CYP3A inducers (ex. carbamazepine, phenobarbital, phenytoin, rifampin, bosentan, dexamethasone, nafcillin, rifabutin, St. John's wort, and others).

Initial approval duration: 6 months

➤ **Criteria for continuation of coverage (renewal request):** Retevmo (selpercatinib) is considered **medically necessary** and will be approved when **ALL** the following criteria are met (**samples are not considered for continuation of therapy**):

1. Individual continues to be seen by a physician specializing in the patient's diagnosis or is in consultation with an Oncologist.
2. Individual's condition has responded while on therapy with response defined as **ALL** of the following:
 - a. Documented evidence of disease stability
 - b. No evidence of disease progression
3. Individual has been adherent with the medication
4. Individual has not developed any significant adverse drug effects that may exclude continued use such as:
 - a. Hepatotoxicity
 - b. Life-threatening hypertension that is not controlled by antihypertensive medications
 - c. Life-threatening QT prolongation
 - d. Severe or life-threatening hemorrhage
 - e. Recurrence of hypersensitivity
5. Individual does not have active cardiovascular disease or a recent myocardial infarction.

ORIGINAL EFFECTIVE DATE: 08/20/2020 | ARCHIVE DATE: | LAST REVIEW DATE: 08/18/2022 | LAST CRITERIA REVISION DATE: 08/18/2022

PHARMACY COVERAGE GUIDELINE

RETEVMO™ (selpercatinib) oral

- 6. Individual does not have end-stage renal disease (eGFR less than 15 mL/min).
- 7. Individual has not had more than 3 dose reductions due to drug toxicity.
- 8. Individual is not using strong and moderate CYP3A inducers (ex. carbamazepine, phenobarbital, phenytoin, rifampin, bosentan, dexamethasone, nafcillin, rifabutin, St. John’s wort, and others).

Renewal duration: 12 months

➤ Criteria for a request for non-FDA use or indication, treatment with dosing, frequency, or duration outside the FDA-approved dosing, frequency, and duration, refer to one of the following Pharmacy Coverage Guideline:

- 1. **Off-Label Use of Non-Cancer Medications**
- 2. **Off-Label Use of Cancer Medications**

Description:

Retevmo (selpercatinib) is indicated for the treatment of adult patients (18 years of age or older) with metastatic *RET* fusion-positive non-small cell lung cancer (NSCLC); for the treatment of adult and pediatric patients 12 years of age and older with advanced or metastatic *RET*-mutant medullary thyroid cancer (MTC) who require systemic therapy; and for the treatment of adult and pediatric patients 12 years of age and older with advanced or metastatic *RET* fusion-positive thyroid cancer who require systemic therapy and who are radioactive iodine-refractory (if radioactive iodine is appropriate).

These indications are approved under accelerated approval based on overall response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s).

Definitions:

Activities of daily living (ADL):

Instrumental ADL:

Prepare meals, shop for groceries or clothes, use the telephone, manage money, etc.

Self-care ADL:

Bathe, dress and undress, feed self, use the toilet, take medications, not bedridden

Common Terminology Criteria for Adverse Events (CTCAE) Version 4.0:

Grade 1	Mild; asymptomatic or mild symptoms; clinical or diagnostic observations only; intervention not indicated
Grade 2	Moderate; minimal, local or noninvasive intervention indicated; limiting age-appropriate instrumental ADL*
Grade 3	Severe or medically significant but not immediately life-threatening; hospitalization or prolongation of hospitalization indicated; disabling; limiting self-care ADL**
Grade 4	Life-threatening consequences; urgent intervention indicated
Grade 5	Death related to AE
U.S. Department of Health and Human Services, National Institutes of Health, and National Cancer Institute	

PHARMACY COVERAGE GUIDELINE

RETEVMO™ (selpercatinib) oral

ECOG Performance status:

Eastern Co-operative Oncology Group (ECOG) Performance Status	
Grade	ECOG description
0	Fully active, able to carry on all pre-disease performance without restriction
1	Restricted in physically strenuous activity but ambulatory and able to carry out work of a light or sedentary nature, e.g., light house work, office work
2	Ambulatory and capable of all selfcare but unable to carry out any work activities. Up and about more than 50% of waking hours
3	Capable of only limited selfcare, confined to bed or chair more than 50% of waking hours
4	Completely disabled. Cannot carry on any self-care. Totally confined to bed or chair
5	Dead

Oken, M.M., Creech, R.H., Tormey, D.C., Horton, J., Davis, T.E., McFadden, E.T., Carbone, P.P.: Toxicity And Response Criteria Of The Eastern Cooperative Oncology Group. Am J Clin Oncol 5:649-655, 1982

Resources:

Retevmo (selpercatinib) product information, revised by Eli Lilly and Company 01-2021. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed July 28, 2021.

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology: Non-Small Cell Lung Cancer. Version 3.2022 – Updated March 16, 2022. Available at <https://www.nccn.org>. Accessed August 03, 2022.

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology: Thyroid Carcinoma. Version 02.2022 – Updated May 05, 2022. Available at <https://www.nccn.org>. Accessed August 03, 2022.

Off Label Use of Cancer Medications: A.R.S. §§ 20-826(R) & (S). Subscription contracts; definitions.

Off Label Use of Cancer Medications: A.R.S. §§ 20-1057(V) & (W). Evidence of coverage by health care service organizations; renewability; definitions.